


1. Prepare for the injection

1a. Remove carton from refrigerator



- Take the carton containing the prefilled syringe out of the refrigerator

1b. Gather your supplies



Cotton ball/gauze and alcohol swab



Adhesive bandage



FDA-cleared sharps disposal container

1c. Let syringe reach room temperature



30-45 minutes


- Allow the prefilled syringe to come to room temperature for 30-45 minutes

1d. Wash and dry your hands



- Wash and dry your hands
- Check your syringe carefully:
 - Do not use if it is damaged or expired
 - Do not use if the medication looks cloudy, discolored, or contains particles

1e. Choose and clean the injection site



X DO NOT inject through:



clothing



moles



scars



bruises




tender skin

If your dose requires more than 1 injection, choose a different injection site for each injection.

2. Give the injection

2a. Remove the syringe cap



- Hold the body of the prefilled syringe and gently remove the cap with your other hand

2b. Dispose of syringe cap




- Dispose of the cap in your sharps container[†]

[†]Do not recap or touch the needle.


3. How to inject

3a. Insert the needle




- Pinch a fold of skin at the injection site and insert the needle into the fold at a 45° angle

3b. Use the correct needle angle for injection



4a. Inject the medicine




- Inject the medication. Keep the skin pinched until you complete the injection

4b. Dispose of the syringe



- Remove the prefilled syringe from the injection site and dispose in your sharps FDA-cleared disposal container

Care for the injection site



- If bleeding occurs, gently press the area with the cotton ball or gauze. Do not rub. Apply the adhesive bandage if needed